

MESSENGER-INQUIRER



May 4, 2024

Some Trees Affected by Anthracnose Causing Leaf Drop
Annette Meyer Heisdorffer, Ph.D.
Extension Agent for Horticulture Education – Daviness County

Green leaves covering the ground in the spring catches our attention. Upon further investigation, the leaves have brown and black patches on them. Looking up in the tree, the brown to black patches are noticeable. Eventually the affected leaves fall off and litter the ground. Anthracnose is a common spring fungal disease causing these blotches. Dr. Nicole Ward, University of Kentucky plant pathologist, provides information about the disease and how to manage it.

The term anthracnose refers to the symptoms of dark blotching and leaf distortion. Leaves drop from the tree during severe infections. The disease is not fatal to the tree but puts stress on it. A new flush of leaves may immediately follow. Twigs and branches infected with the fungi develop cankers and girdled stems.

Anthracnose diseases on different species of trees are not caused by the same fungi. The pathogens that cause the disease are host-specific which means the anthracnose pathogen on

MESSENGER-INQUIRER

sycamore will not infect ash and so on. The appearance of the blackened area and severity of disease differs with each host and weather conditions.

Even though different, the fungal pathogens that cause anthracnose diseases have similar life cycles. Spore production occurs during periods of spring rain. Anthracnose fungi release spores to infect deciduous trees when leaves emerge. Mature leaves are resistant to infection, but slow leaf emergence in spring exposes tender leaves to fungal spores for an extended period of time.

Ash anthracnose, caused by *Discula umbrinella*, infects buds, leaves, and sometimes twigs. In early spring, infection of buds or expanding leaves results in irregular brown blotches and leaflet distortion. These blotches are frequently associated with leaf margins. Infections that occur once leaves have already expanded result in small brown circular lesions. As these lesions enlarge, they may come together forming one big brown area. Infected leaflets often drop. Although shoots may become stunted, infection on ash does not result in obvious twig or branch cankers.

Maple anthracnose symptoms begin as leaf spots with dead tissue and may progress into shoot blight and shoot cankers. The way the disease looks on the leaves varies in size and shape depending on the maple species. At least two different anthracnose fungi may be involved, *Discula* sp. and *Kabatiella apocrypta*. On Norway maple, lesions or spots are purple to brown and follow the veins. Leaves of Japanese maple blacken and shrivel up. Brown to reddish brown lesions form along or between veins of sugar maple. Symptoms on sugar maple can be confused with scorch symptoms. Twig infections result in blighting and death of shoot tips.

MESSENGER-INQUIRER

Sycamore anthracnose on young, expanding leaves develops irregular dark, necrotic blotching centered along leaf veins or edges. These dark blotches may turn tan-colored as the diseased areas of the leaves dry out. Blighting of twigs or shoots may follow. Trees produce new foliage rather quickly, but affected branches may remain crooked because lateral shoots behind the terminal or end shoot, become dominant when terminal shoots are killed.

Besides the foliar infection, persistent rains and disease spread can lead to infection of twigs and branches. Good cultural practices, like those below, are important to manage this fungal disease.

Anthracnose is favored by a moist environment. Select a planting site with a sunny eastern exposure to promote rapid foliage drying early in the day.

Remember to rake and destroy fallen leaves, as they can be a source of inoculum or fungal spores. Do not compost infected leaves. Remove dead twigs and branches which serve as an overwintering site for the fungi.

Avoid wounding tree trunks, such as bumping with lawn equipment and making jagged pruning cuts.

Maintain mulch 2-3 inches thick over the root zone and beyond the drip line but not against the trunk to maintain soil moisture and protect trees from lawnmower injury.

Protect trees from dry weather and drought. Water the trees at least once a week during hot, dry months using soaker hoses or drip irrigation. Avoid overhead sprinklers; wet foliage favors sporulation and infection.

Diagnose insect and disease problems as soon as possible and treat if appropriate.

MESSENGER-INQUIRER

Fungicides are often not recommended to manage anthracnose. They can be costly, and it is difficult to effectively cover large trees. These chemicals are protectants and therefore must be applied before infection occurs. Once symptoms develop, it is too late to apply fungicides for controlling anthracnose. However, if the tree is valuable or if it has been attacked year after year, a fungicide spray program may be justified. Three sprays should be applied in spring: at bud break, when leaves are half-expanded, and when leaves are fully expanded. Good coverage with the fungicide is needed.

For more information about anthracnose on shade trees, contact the Daviess County Cooperative Extension Service at 270-685-8480 or annette.heisdorffer@uky.edu.

Annette's Tip:

When looking for information about growing plants in the garden and landscape, consider looking for information through the University of Kentucky at Daviess.ca.uky.edu. On this site, click on horticulture. A variety of topics and links for resources are presented.

Also, the past gardening programs I have presented at the Daviess County Public Library on gardening are available through their Facebook page.

On Going Event:

The state-wide Horticulture Webinar Wednesday is offered from 11:30 a.m. to noon. "Home Cut Flower Gardening" is the next topic. The site to register is located at <https://kentuckyhortnews.com/>. Presentations are recorded and topics are found there for the past four years.

MESSENGER-INQUIRER

The Martin-Gatton College of Agriculture, Food and Environment is an Equal Opportunity Organization with respect to education and employment and authorization to provide research, education information and other services only to individuals and institutions that function without regard to economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, physical or mental disability or reprisal or retaliation for prior civil rights activity.